

Advanced Quantitative Research Methodology, Lecture Notes: Theories of Inference¹

Gary King
<http://GKing.Harvard.Edu>

February 25, 2013

¹©Copyright 2013 Gary King, All Rights Reserved.

The Problem of Inference

1. Probability: $P(y|M) = P(\text{known}|\text{unknown})$
2. The goal of inverse probability: $P(M|y) = P(\text{unknown}|\text{known})$.
3. Define: $M = \{M^*, \theta\}$, where M^* is assumed and θ is to be estimated
4. More limited goal: $P(\theta|y, M^*) \equiv P(\theta|y)$.

The Problem of Inference

5. Bayes Theorem (no additional assumptions):

$$\begin{aligned} P(\theta|y) &= \frac{P(\theta, y)}{P(y)} && [\text{Defn. of conditional probability}] \\ &= \frac{P(\theta)P(y|\theta)}{P(y)} && [P(AB) = P(B)P(A|B)] \\ &= \frac{P(\theta)P(y|\theta)}{\int P(\theta)P(y|\theta)d\theta} && [P(A) = \int P(AB)dB] \end{aligned}$$

6. If we knew the right side, we could compute the inverse probability.
7. 2 theories of inference arose to interpret this result: likelihood and Bayesian
8. In both, $P(y|\theta)$ is a traditional probability density
9. The two differ on the rest

Interpretation 1: The Likelihood Theory of Inference

1. R.A. Fisher's idea
2. θ is fixed and y is random
3. Let:

$$k(y) \equiv \frac{P(\theta)}{\int P(\theta)P(y|\theta)d\theta} \implies P(\theta|y) = \frac{P(\theta)P(y|\theta)}{\int P(\theta)P(y|\theta)d\theta} = k(y)P(y|\theta)$$

4. Define $K(y)$ as an *unknown* function of y with θ fixed at its true value
5. $\rightsquigarrow$ the likelihood theory of inference has four axioms: the 3 probability axioms plus the **likelihood axiom**:

$$\begin{aligned} L(\theta|y) &\equiv k(y)P(y|\theta) \\ &\propto P(y|\theta) \end{aligned}$$


6. $L(\theta|y)$ is a function: for y fixed at the observed values, it gives the “likelihood” of any value of θ .

Interpretation 1: The Likelihood Theory of Inference

7. Likelihood: a **relative measure of uncertainty**, changing with the data set
8. Comparing the value of $L(\theta|y)$ for different θ values in one data set y is meaningful.
9. Comparing values of $L(\theta|y)$ across data sets is meaningless. (just as you can't compare R^2 values across equations with different dependent variables.)
10. The **likelihood principle**: the data only affect inferences through the likelihood function

Visualizing the Likelihood

- For algebraic simplicity and numerical stability, we use the **log-likelihood** (the shape changes, but the max is in the same place)
- If θ has one element, we can plot:



- The full likelihood curve is a **Summary Estimator**. The likelihood principle means that once this is plotted, we can discard the data (if the model is correct!).
- A one-point summary at the maximum is the **MLE**
- Uncertainty of point estimate: **curvature at the maximum**

Interpretation 2: The Bayesian Theory of Inference

- Rev. Thomas Bayes' unpublished idea, and later rediscovered.
- Recall:

$$\begin{aligned} P(\theta|y) &= \frac{P(\theta, y)}{P(y)} && [\text{Defn. of conditional probability}] \\ &= \frac{P(\theta)P(y|\theta)}{P(y)} && [P(AB) = P(B)P(A|B)] \\ &= \frac{P(\theta)P(y|\theta)}{\int P(\theta)P(y|\theta)d\theta} && [P(A) = \int P(AB)dB] \\ &\propto P(\theta)P(y|\theta) \end{aligned}$$

- $P(\theta|y)$ the posterior density
- $P(y|\theta)$ the traditional probability ($\propto$ likelihood)
- $P(y)$ a constant, easily computed
- $P(\theta)$, the prior density — the way Bayes differs from likelihood


What is the prior density, $P(\theta)$?

1. A probability density that represents all prior evidence about θ .
2. An opportunity: a way of getting other information outside the data set into the model
3. An annoyance: the “other information” is required
4. A philosophical assumption that nonsample information should matter (as it always does) *and* be formalized and included in all inferences.

Principles of Bayesian analysis

1. All unknown quantities (θ, Y) are treated as random variables and have a joint probability distribution.
2. All known quantities (y) are treated as fixed.
3. If we have observed variable B and unobserved variable A , then we are usually interested in the conditional distribution of A , given B :
$$P(A|B) = P(A, B)/P(B)$$
4. If variables A and B are both unknown, then the distribution of A alone is $P(A) = \int P(A, B)dB = \int P(A|B)P(B)dB$.

The posterior density, $P(\theta|y)$



- Like L , it's a summary estimator
- Unlike L , it's a real probability density, from which we can derive probabilistic statements (via integration)
- To compare across applications or data sets, you may need different priors. So, the posterior is also relative, just like likelihood.
- **Bayesian inference** obeys the **likelihood principle**: the data set only affects inferences through the likelihood function
- If $P(\theta) = 1$, i.e., is uniform in the relevant region, then $L(\theta|y) = P(\theta|y)$.

Bayesians are happier people

- If $P(\theta)$ is *diffuse*, differences from likelihood are minor, but numerical stability (and “identification”) is improved (your programs will run better!).
- Philosophical differences from likelihood: **Huge**
- Practical differences when we can compute both: **Minor** (unless the prior matters)
- Advantages: more information produces more efficiency; MCMC algorithms are easier with Bayes.
- Few fights now between Bayesians and likelihoodists

A 3rd Theory: Neyman-Pearson Hypothesis Testing

1. Huge fights between these folks and the {Bayesians, Likelihoodists}
2. Strict but arbitrary distinction: null H_0 vs alternative H_1 hypotheses
3. All tests are “under” (i.e., assuming) H_0
4. For example, is $\beta = 0$ in $E(Y) = \beta_0 + \beta X$?
 - (a) $H_0: \beta = 0$ vs. $H_1: \beta > 0$
 - (b) Choose Type I error, probability of deciding H_1 is right when H_0 is really true: say $\alpha = 0.05$
 - (c) Type II error, the power to detect H_1 if it is true, is a consequence of choosing an estimator, not an ex ante decision like choosing α .
 - (d) Assume n is large enough for the CLT to kick in
 - (e) Then $b|(\beta = 0) \sim N(0, \sigma_b^2)$
 - (f) or

$$(TS)_{\beta}|(\beta = 0) \equiv \frac{b - \beta}{\hat{\sigma}_b} \equiv \frac{b}{\hat{\sigma}_b} \sim N(0, 1).$$

Neyman-Pearson Hypothesis Testing

- (g) Derive critical value, CV , e.g., the right tail:


$$\int_{(CV)}^{\infty} N(b|0, \sigma_b^2) db = \alpha$$

- (h) This means, in educational psychology and other fields, write your prospectus, plan your experiment, report the CV , and write your concluding chapter:

$$\text{Decision} = \begin{cases} \beta > 0 \text{ (I was right)} & \text{if } (TS) > (CV) \\ \beta = 0 \text{ (I was wrong)} & \text{if } (TS) \leq (CV) \end{cases}$$

And then first collect your data. You may not revise your hypothesis or your theory.

Neyman-Pearson Hypothesis Testing



- (i) In this example, $(TS) < (CV)$ and so we conclude that $\beta = 0$.
 - (j) Decision will be wrong 5% of the time; what about this time?
5. What about when n is large or under control of the investigator?
 6. In practice, hypothesis testing is used with p -values: **The probability under the null of getting a value as weird or weirder than the value we got** — the area to the right of the realized value of (TS) .
 7. Star-gazing is silly; where is the quantity of interest?
 8. We can use likelihood to compute hypothesis tests and p -values.

What is the right theory of inference?

1. Likelihood? Bayes? Neyman-Pearson? Criteria estimators? Finite or asymptotic based theory? Decision theory? Nonparametrics? Semiparametrics? Conditional inference? Superpopulation-based inference? etc.
2. None of these.
3. The right theory of inference: **utilitarianism**

Unification of Theories of Inference

- Can't bank on agreement on normative issues!
- Even if there is agreement, it won't hold or shouldn't
- Alternative convergence is occurring: different methods giving the same result.
 - Likelihood or Bayes with careful goodness of fit checks
 - Various types of robust or semi-parametric methods
 - Matching for use as preprocessing for parametric analysis
 - Bayesian model averaging, with a large enough class of models to average over
 - Committee methods, mixture of experts models
 - Some models with highly flexible functional forms
- The key: No assumptions can be trusted; all theories of inference condition on assumptions and so data analysts always struggle trying to understand and get around them

A Simple Likelihood Model: Stylized Normal, no X

The model:

1. $Y_i \sim f_{\text{stn}}(y_i|\mu_i)$, normal **stochastic** component
2. $\mu_i = \beta$, a constant **systematic** component (no covariates)
3. Y_i and Y_j are independent $\forall i \neq j$.

Derive the full probability density of *all* observations, $\Pr(\text{data}|\text{model})$
(Recall: if A and B are independent, $P(AB) = P(A)P(B)$):

$$\begin{aligned} P(y|\mu) &\equiv P(y_1, \dots, y_n|\mu_1, \dots, \mu_n) = \prod_{i=1}^n f_{\text{stn}}(y_i|\mu_i) \\ &= \prod_{i=1}^n (2\pi)^{-1/2} \exp\left(\frac{-(y_i - \mu_i)^2}{2}\right) \end{aligned}$$

reparameterizing with $\mu_i = \beta$:

$$P(y|\beta) \equiv P(y_1, \dots, y_n|\beta) = \prod_{i=1}^n (2\pi)^{-1/2} \exp\left(\frac{-(y_i - \beta)^2}{2}\right)$$

- What can you do with this density?

Stylized Normal Likelihood Function


$$\begin{aligned}L(\beta|y) &= k(y) \prod_{i=1}^n f_{\text{stn}}(y_i|\beta) \\&\propto \prod_{i=1}^n f_{\text{stn}}(y_i|\beta) \\&= \prod_{i=1}^n (2\pi)^{-1/2} \exp\left(\frac{-(y_i - \beta)^2}{2}\right)\end{aligned}$$

the relative likelihood of β (conditional on the model) having generated the data we observe.

The log-likelihood (Recall: $\ln(ab) = \ln(a) + \ln(b)$):

$$\begin{aligned}\ln L(\beta|y) &= \ln[k(y)] + \sum_{i=1}^n \ln f_{\text{stn}}(y_i|\beta) \\&= \ln[k(y)] + \sum_{i=1}^n \ln[(2\pi)^{-1/2}] - \sum_{i=1}^n \frac{1}{2}(y_i - \beta)^2 \\&\doteq \sum_{i=1}^n -\frac{1}{2}(y_i - \beta)^2 \\&= -\frac{1}{2} \sum_{i=1}^n (y_i - \beta)^2\end{aligned}$$

Log-likelihood interpretation



1. The log-likelihood is quadratic
2. The MLE is at the same point as the MVLUE
3. The maximum is at the same point as the least squares point
4. By the theory, this curve summarizes all information the data gives about β , assuming the model.
5. No reason to summarize this curve with only the MLE

Summarizing k -dimensional space

- The problem of Flatland
- Graphs
- Maximum
- The curvature at the maximum (standard errors, about which more shortly)
- The curse of dimensionality

How to find the maximum?

Goal: Find the value of $\theta \equiv \{\theta_1, \dots, \theta_k\}$ that maximizes $L(\theta|y)$

① Analytically — often impossible or too hard

- Take the derivative of $\ln L(\theta|y)$ w.r.t. θ
- Set to 0, substituting $\hat{\theta}$ for θ

$$\left| \frac{\partial \ln L(\theta|y)}{\partial \theta} \right|_{\theta=\hat{\theta}} = 0$$

- If possible, solve for θ , and label it $\hat{\theta}$
- Check the second order condition: see if the second derivative w.r.t. θ is negative (so its a maximum rather than a minimum)

② Numerically — let the computer do the work for you

- We'll show you how

Finite Sample Properties of the MLE

1 Minimum variance unbiased estimator (MVUE)

- Unbiasedness:
 - Definition: $E(\hat{\theta}) = \theta$
 - Example: $E(\bar{Y}) = E\left(\frac{1}{n} \sum_{i=1}^n Y_i\right) = \frac{1}{n} \sum_{i=1}^n E(Y_i) = \frac{1}{n} n\mu = \mu$
- Minimum variance (“efficiency”)
 - Variance to be minimized: $V(\hat{\theta})$
 - Example: $V(\bar{Y}) = V\left(\frac{1}{n} \sum_{i=1}^n Y_i\right) = \frac{1}{n^2} \sum_{i=1}^n V(Y_i) = \frac{1}{n^2} n\sigma^2 = \sigma^2/n$
- If there is a MVUE, ML will find it
- If there isn't one, ML will still usually find a good estimator, even when other methods of inference won't

2 Invariance to Reparameterization

- Estimate σ with $\hat{\sigma}$ and calculate $\hat{\sigma}^2$ or estimate $\hat{\sigma}^2$: both are MLEs
- Not true for other methods of inference: e.g. $\bar{y}$ is an unbiased estimate of μ . What is an unbiased estimate of $1/\mu$? $E(1/\bar{y}) \neq 1/E(\bar{y})$.


3 Invariance to sampling plans

- OK to look at results while deciding how much data to collect
- In fact, it's a great idea!


Asymptotic Properties of the MLE

- 1 **Consistency** (from the Law of Large Numbers). As $n \rightarrow \infty$, the sampling distribution of the MLE collapses to a spike over the parameter value
- 2 **Asymptotic normality** (from the central limit theorem). As $n \rightarrow \infty$, the distribution of $\text{MLE}/\text{se}(\text{MLE})$ converges to a Normal.
 - Do the LLN and CLT (the 2 most important theorems in statistics) contradict each other?
- 3 **Asymptotic efficiency**. The MLE contains as much information as can be packed into a point estimator.

Sampling distributions of the MLE: CLT vs LLN



Uncertainty: Likelihood Ratios for nested models



- L^* is the likelihood value for the **unrestricted** model
- L_R^* is the likelihood value for the (nested) **restricted** model
- $\implies L^* \geq L_R^* \implies \frac{L_R^*}{L^*} \leq 1$

Meaning of the likelihood ratio

1. Substantively, its the ratio of 2 traditional probabilities:

$$\begin{aligned}L(\theta_1|y) &\propto k(y)P(y|\theta_1) \\L(\theta_2|y) &\propto k(y)P(y|\theta_2) \\ \frac{L(\theta_1|y)}{L(\theta_2|y)} &= \frac{k(y)}{k(y)} \frac{P(y|\theta_1)}{P(y|\theta_2)} \\ &= \frac{P(y|\theta_1)}{P(y|\theta_2)}\end{aligned}$$

Interpreted as a *risk ratio*.

2. Statistically (from the Neyman-Pearson Hypothesis Testing viewpoint), let

$$R = -2 \ln \left(\frac{L_R^*}{L^*} \right) = 2(\ln L^* - \ln L_R^*)$$


Then, under the null of no difference between the 2 models,

$$R \sim f_{\chi^2}(r|m)$$

where r is the observed value of R and m is the number of restricted parameters.


Meaning of the likelihood ratio

- (a) If restrictions have no effect, $E(R) = m$.
- (b) So only if $r \gg m$ will the test parameters be clearly different from zero.



- 3. Disadvantage: Too many likelihood ratio tests may be required to test all points of interest

Standard Errors



1. Instead of (a) plotting the entire likelihood hyper-surface or (b) computing numerous likelihood ratio tests, we can summarize all the curvature near the maximum with one number.
2. We will use the normal likelihood to approximate all likelihoods
3. (one justification) as $n \rightarrow \infty$, likelihoods become normal.
4. Reformulate the normal (not stylized normal) likelihood.

$$\begin{aligned} L(\beta|y) &\propto N(y_i|\mu_i, \sigma^2) \\ &= (2\pi\sigma^2)^{-1/2} \exp\left(\frac{-(y_i - \mu_i)^2}{2\sigma^2}\right) \\ &= (2\pi\sigma^2)^{-1/2} \exp\left(\frac{-(y_i - \beta)^2}{2\sigma^2}\right) \end{aligned}$$

Justifying Standard Errors

$$\begin{aligned}\ln L(\beta|y) &= -\frac{n}{2} \ln(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \beta)^2 \\ &= -\frac{n}{2} \ln(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i^2 - 2y_i\beta + \beta^2) \\ &= \left(-\frac{n}{2} \ln(2\pi\sigma^2) - \frac{\sum_{i=1}^n y_i^2}{2\sigma^2} \right) + \left(\frac{\sum_{i=1}^n y_i}{\sigma^2} \right) \beta + \left(\frac{-n}{2\sigma^2} \right) \beta^2 \\ &= a + b\beta + c\beta^2, \quad \text{A quadratic equation}\end{aligned}$$

5. $\left(\frac{-n}{2\sigma^2}\right)$ is the degree of curvature. Curvature is larger when:
- n is large
 - σ^2 is small
6. For normal likelihood, $\left(\frac{-n}{2\sigma^2}\right)$ is a summary. The bigger the (negative) number...
- the better
 - the more information exists in the MLE
 - the larger the likelihood ratio would be in comparing the MLE with *any* other parameter value.

Standard Errors

7. When the log-likelihood is not normal, we'll use the best quadratic approximation to it. Under the normal,

$$\frac{\partial^2 \ln L(\beta|y)}{\partial \beta \partial \beta'} = \frac{-n}{\sigma^2}$$

More generally, this second derivative will give us a way to compute the coefficient on the squared term.

8. We invert the curvature to provide a statistical interpretation:

$$\hat{V}(\hat{\theta}) = \left[-\frac{\partial^2 \ln L(\theta|y)}{\partial \theta \partial \theta'} \right]_{\theta=\hat{\theta}}^{-1} = \begin{pmatrix} \hat{\sigma}_1^2 & \hat{\sigma}_{12} & \cdots \\ \hat{\sigma}_{21} & \hat{\sigma}_2^2 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix}$$

- Statistical interpretation: variance and covariance across repeated samples
- Works in general for a k -dimensional θ vector
- Can be computed numerically
- Known as the variance matrix, or variance-covariance matrix, or covariance matrix

9. This is an **estimate** of a **quadratic approximation** to the log-likelihood.
10. Asymptotically, it is an estimate of the exact log-likelihood.

Simulation for any ML Model

- If the model is correct, a consistent point estimate of θ is the MLE, $\hat{\theta}$.
- As n gets large,
 - the sampling distribution of $\hat{\theta}$ becomes normal.
 - the log-likelihood becomes quadratic
 - the quadratic approximation assumed when using the second derivative of the log-likelihood improves
- Estimate of the variance of the sampling distribution of $\hat{\theta}$: $\hat{V}(\hat{\theta})$, the inverse of the negative of the matrix of second derivatives, evaluated at $\hat{\theta}$.
- To simulate θ ,
 - we'll draw from the multivariate normal: $N(\hat{\theta}, \hat{V}(\hat{\theta}))$
 - This is an asymptotic approximation and can be wrong sometimes.
 - We'll discuss later how to improve the approximation.

ML Example: k Parameters, including an Ancillary Parameter, with Simulation to Interpret.

Forecasting Presidential Elections.

The Data

i	U.S. state, for $i = 1, \dots, 50$
t	election year, for $t = 1948, 1952, \dots, 2008$
y_{it}	Democratic fraction of the two-party vote
X_{it}	a list of covariates (economic conditions, polls, home state, etc)
$X_{i,2012}$	the same covariates as X_{it} but measured in 2012
E_i	The number of electoral college votes for each state in 2012

The Model

1. $Y_{it} \sim N(\mu_{it}, \sigma^2)$.
2. $\mu_{it} = x_{it}\beta$, where x_{it} is a vector of explanatory variables and a constant
3. Y_{it} and $Y_{i't'}$ are independent $\forall i \neq i'$ and $t \neq t'$, conditional on X .

The Likelihood Model for the i th observation

$$\begin{aligned} L(\mu_{it}, \sigma | y_{it}) &\propto N(y_{it} | \mu_{it}, \sigma^2) \\ &= (2\pi\sigma^2)^{-1/2} e^{\frac{-(y_{it} - \mu_{it})^2}{2\sigma^2}} \end{aligned}$$

Likelihood model for all n observations

$$\begin{aligned}L(\beta, \sigma^2 | y) &= \prod_{i=1}^n \prod_{t=1}^T L(y_{it} | \mu_{it}, \sigma^2) \\ \ln L(\beta, \sigma^2 | y) &= \sum_{i=1}^n \sum_{t=1}^T \ln L(y_{it} | \mu_{it}, \sigma^2) \\ &= \sum_{i=1}^n \sum_{t=1}^T \left\{ -\frac{1}{2} \ln(2\pi\sigma^2) - \frac{(y_{it} - \mu_{it})^2}{2\sigma^2} \right\} \\ &= \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \ln(2\pi) + \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \left[\ln \sigma^2 + \frac{(y_{it} - \mu_{it})^2}{\sigma^2} \right] \\ &= \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \left[\ln \sigma^2 + \frac{(y_{it} - \mu_{it})^2}{\sigma^2} \right] \\ &= \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \left[\ln \sigma^2 + \frac{(y_{it} - \mathbf{X}_{it}\beta)^2}{\sigma^2} \right]\end{aligned}$$

- k : number of explanatory variables
- Reparameterize on the unbounded scale; use: $\sigma = e^\gamma$
- Let $\theta = \{\beta, \gamma\}$, a $k + 2 \times 1$ vector.
- Maximize the likelihood; save $\hat{\theta} = \{\hat{\beta}, \hat{\gamma}\}$.
- Compute and save $\hat{V}(\hat{\theta})$, which is $k + 2 \times k + 2$

R Code for the Log-Likelihood

- Mathematical Form:

$$\ln L(\beta, \sigma^2 | y) = \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \left[\ln \sigma^2 + \frac{(y_{it} - X_{it}\beta)^2}{\sigma^2} \right]$$

- The function:

```
ll.normal <- function(par, X, Y) {  
  beta <- par[1:ncol(X)]  
  sigma2 <- exp(par[ncol(X) + 1])  
  -1/2 * sum( log(sigma2) + ((Y - X %*% beta)^2)/sigma2 )  
}
```

- Calling it:

```
ll.normal(c(2,1,2,1,33,4,3.2),x,y)  
ll.normal(c(2,1,2,1,33,4,3.7),x,y)  
ll.normal(c(2,1,2,1,33,4,3.5),x,y)
```

Quantities of Interest

- Reasons we care about the regression coefficients: None
- The posterior distribution of electoral college delegates for the Democrat.
- Expected number of electoral college delegates for the Democrat.
- Probability that the Democratic candidate gets more than $\sum_{i=1}^n E_i / n > 0.5$ proportion of electoral college delegates.

Predictive distribution of electoral college delegates in 2012

- Draw many simulations of $y_{i,2012}$ ($\tilde{y}_{i,2012}$) from its posterior distribution for U.S. state i , $P(y_{i,2012}|y_{it}, X_{it})$, i.e. $P(\text{quantity of interest}|\text{data})$. (Details shortly.)
- For each simulation of state i , if $y_{i,2012} > 0.5$ the Democrat “wins” E_i electoral college delegates; if its not, he gets zero.
- Add the number of electoral college delegates the Democrat wins in the entire country by adding simulated winnings from each state.
- Repeat Steps 1–3 $M = 1,000$ times, and plot a histogram of the results.
- (How does this differ from allocating all E_i based on the point estimate?)


How to draw simulations of $y_{i,2012}$?

1. Choose values of explanatory variables. In this case, $X_{i,2012}$
2. Simulate estimation uncertainty:
 - Draw θ randomly from its posterior (i.e., sampling) distribution, $N(\theta|\hat{\theta}, \hat{V}(\hat{\theta}))$. Label the random draw $\tilde{\theta} = \{\hat{\beta}, \hat{\gamma}\}$.
 - Pull out $\tilde{\beta}$ and save.
 - Pull out, un-reparameterize, and save $\tilde{\sigma} = e^{\tilde{\gamma}}$
3. Compute the simulated systematic component: $\tilde{\mu}_{it} = X_{i,2012}\tilde{\beta}$
4. Add fundamental uncertainty: draw $y_{i,2012}$ from $N(y_{i,2012}|\tilde{\mu}_{i,2012}, \tilde{\sigma}^2)$

How to do it with a LS Regression Program

1. Run LS regression of y_{it} on X_{it} and get $\hat{\beta}$ and $V(\hat{\beta})$
2. Draw β randomly from its posterior distribution (i.e., its sampling distribution), $N(\beta|\hat{\beta}, V(\hat{\beta}))$. Label the random draw $\tilde{\beta}$.
3. Draw σ^2 from its posterior (or sampling) distribution, $1/\chi^2(\sigma^2|\hat{\sigma}^2, N - k)$, labeling it $\tilde{\sigma}^2$
4. Either:
 - Draw ϵ_{it} from $N(\epsilon_{it}|0, \tilde{\sigma}^2)$, label it $\tilde{\epsilon}_{it}$ and compute:
$$\tilde{y}_{i,2012} = \tilde{X}_{i,2012}\tilde{\beta} + \tilde{\epsilon}_{it}$$
 - Or, in our preferred notation, draw $\tilde{y}_{i,2012}$ from $N(X_{i,2012}\tilde{\beta}, \tilde{\sigma}^2)$

Actual Results (calculated before the election) for 1992



Variance Function Models


1. $Y_{it} \sim N(y_{it}|\mu_{it}, \sigma_{it}^2)$
2. $\mu_{it} = x_{it}\beta$, where x_{it} is a vector of explanatory variables and a constant
3. $\sigma_{it}^2 = \exp(z_{it}\gamma)$, where z_{it} is a vector of explanatory variables possibly overlapping x_{it}
4. Y_{it} and $Y_{i't'}$ are independent $\forall i \neq i'$ and $t \neq t'$, conditional on X and Z .

The log-likelihood:

$$\begin{aligned}\ln L(\beta, \sigma^2|y) &= \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \left[\ln \sigma^2 + \frac{(y_{it} - X_{it}\beta)^2}{\sigma^2} \right] \\ &= \sum_{i=1}^n \sum_{t=1}^T -\frac{1}{2} \left[z_{it}\gamma + \frac{(y_{it} - X_{it}\beta)^2}{\exp(z_{it}\gamma)} \right]\end{aligned}$$

- For what applications would this model be informative?

An Outline of the Research Process



1. These figures are always wild simplifications.
2. Items are roughly in order.
3. You can start at any point.
4. A formal theory is often a useful addition, but not always necessary.
5. Don't miss any parts.