

Advanced Quantitative Research Methodology, Lecture Notes: Research Designs for Causal Inference¹

Gary King

GaryKing.org

April 14, 2013

¹©Copyright 2013 Gary King, All Rights Reserved.

- Kosuke Imai, Gary King, and Elizabeth Stuart. **Misunderstandings among Experimentalists and Observationalists: Balance Test Fallacies in Causal Inference** *Journal of the Royal Statistical Society, Series A* Vol. 171, Part 2 (2008): Pp. 1-22
<http://gking.harvard.edu/files/abs/matchse-abs.shtml>

- **Sample** of n units from a finite population of N units (typically $N \gg n$)
- **Sample selection:** I_i is 1 for units selected, 0 otherwise
- **Treatment assignment:** T_i is 1 for treated group, 0 control group
- (**Assume:** treated and control groups are each of size $n/2$)
- **Observed outcome variable:** Y_i
- **Potential outcomes:** $Y_i(1)$ and $Y_i(0)$, Y_i potential values when T_i is 1 or 0 respectively.
- **Fundamental problem of causal inference.** Only one potential outcome is ever observed:
If $T_i = 0$, $Y_i(0) = Y_i$ $Y_i(1) = ?$
If $T_i = 1$, $Y_i(0) = ?$ $Y_i(1) = Y_i$
- (I_i, T_i, Y_i) are random; $Y_i(1)$ and $Y_i(0)$ are fixed.

Quantities of Interest

- Treatment Effect (for unit i):

$$\text{TE}_i \equiv Y_i(1) - Y_i(0)$$

- Population Average Treatment Effect:

$$\text{PATE} \equiv \frac{1}{N} \sum_{i=1}^N \text{TE}_i$$

- Sample Average Treatment Effect:

$$\text{SATE} \equiv \frac{1}{n} \sum_{i \in \{I_i=1\}} \text{TE}_i$$

Decomposition of Causal Effect Estimation Error

- Difference in means estimator:

$$D \equiv \left(\frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} Y_i \right) - \left(\frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=0\}} Y_i \right).$$

- Estimation Error:

$$\Delta \equiv \text{PATE} - D$$

- Pretreatment confounders: X are observed and U are unobserved
- Decomposition:

$$\begin{aligned} \Delta &= \Delta_S + \Delta_T \\ &= (\Delta_{S_X} + \Delta_{S_U}) + (\Delta_{T_X} + \Delta_{T_U}) \end{aligned}$$

Error due to Δ_S (sample selection), Δ_T (treatment imbalance), and each due to observed (X_i) and unobserved (U_i) covariates

- Definition:

$$\begin{aligned}\Delta_S &\equiv \text{PATE} - \text{SATE} \\ &= \frac{N - n}{N}(\text{NATE} - \text{SATE}),\end{aligned}$$

where NATE is the nonsample average treatment effect.

- Δ_S vanishes if:
 - 1 The sample is a census ($I_i = 1$ for all observations and $n = N$);
 - 2 $\text{SATE} = \text{NATE}$; or
 - 3 Switch quantity of interest from PATE to SATE (recommended!)

Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_X} + \Delta_{S_U}$$

- $\Delta_{S_X} = 0$ when empirical distribution of (observed) X is identical in population and sample: $\tilde{F}(X | I = 0) = \tilde{F}(X | I = 1)$.
- $\Delta_{S_U} = 0$ when empirical distribution of (unobserved) U is identical in population and sample: $\tilde{F}(U | I = 0) = \tilde{F}(U | I = 1)$.
- conditions are unverifiable: X is observed only in sample and U is not observed at all.
- Δ_{S_X} vanishes if weighting on X
- Δ_{S_U} cannot be corrected after the fact

Decomposing Treatment Imbalance

- Decomposition:

$$\Delta_T = \Delta_{T_X} + \Delta_{T_U}$$

- $\Delta_{T_X} = 0$ when X is balanced between treateds and controls:

$$\tilde{F}(X \mid T = 1, I = 1) = \tilde{F}(X \mid T = 0, I = 1).$$

Verifiable from data; can be generated ex ante by blocking or enforced ex post via matching or parametric adjustment

- $\Delta_{T_U} = 0$ when U is balanced between treateds and controls:

$$\tilde{F}(U \mid T = 1, I = 1) = \tilde{F}(U \mid T = 0, I = 1).$$

Unverifiable. Achieved only by assumption or, on average, by random treatment assignment

Alternative Quantities of Interest

- Sample average treatment effect on the treated:

$$\text{SATT} \equiv \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} \text{TE}_i$$

- Population average treatment effect on the treated:

$$\text{PATT} \equiv \frac{1}{N^*} \sum_{i \in \{T_i=1\}} \text{TE}_i$$

(where $N^* = \sum_{i=1}^N T_i$ is the number of treated units in the population)

- When are these of more interest than PATE and SATE? Why never for randomized experiments? Why usually for matching?
- Analogous estimation error decomposition: $\Delta' = \text{PATT} - D$, holds:

$$\Delta' = (\Delta'_{S_X} + \Delta'_{S_U}) + (\Delta'_{T_X} + \Delta'_{T_U})$$

Effects of Design Components on Estimation Error

Design Choice

	Δ_{S_X}	Δ_{S_U}	Δ_{T_X}	Δ_{T_U}
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$
Exact matching			$= 0$	$= ?$

Assumption

No selection bias	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Ignorability				$\stackrel{\text{avg}}{=} 0$
No omitted variables				$= 0$

Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like parametric adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:
 - ① to avoid selection error, must change quantity of interest from PATE to PATT or SATT,
 - ② random treatment assignment following matching is impossible,
 - ③ Exact matching, unlike blocking is dependent on the already-collected data happening to contain sufficiently good matches.
 - ④ In the worst case scenerio, matching (just like parametric adjustment) can increase bias (this cannot occur with blocking plus random assignment)
- Adding matching to a parametric model almost always reduces model dependence and bias, and sometimes variance too

The Benefits of Major Research Designs: Overview

	Δ_{S_X}	Δ_{S_U}	Δ_{T_X}	Δ_{T_U}
Ideal experiment	$\rightarrow 0$	$\rightarrow 0$	$= 0$	$\rightarrow 0$
Randomized clinical trials (Limited or no blocking)	$\neq 0$	$\neq 0$	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Randomized clinical trials (Full blocking)	$\neq 0$	$\neq 0$	$= 0$	$\stackrel{\text{avg}}{=} 0$
Social Science Field Experiment (Limited or no blocking)	$\neq 0$	$\neq 0$	$\rightarrow 0$	$\rightarrow 0$
Survey Experiment (Limited or no blocking)	$\rightarrow 0$	$\rightarrow 0$	$\rightarrow 0$	$\rightarrow 0$
Observational Study (Representative data set, Well-matched)	≈ 0	≈ 0	≈ 0	$\neq 0$
Observational Study (Unrepresentative but partially, correctable data, well-matched)	≈ 0	$\neq 0$	≈ 0	$\neq 0$
Observational Study (Unrepresentative data set, Well-matched)	$\neq 0$	$\neq 0$	≈ 0	$\neq 0$

For column Q , “ $\rightarrow 0$ ” denotes $E(Q) = 0$ and $\lim_{n \rightarrow \infty} \text{Var}(Q) = 0$, whereas “ $\stackrel{\text{avg}}{=} 0$ ” indicates zero on average, or $E(Q) = 0$, for a design with a small n . Δ_S can be set to zero if we switch from PATE to SATE.

The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large n
- blocking on all known confounders
- random treatment assignment within blocks
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $\Delta_{T_X} = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large n
- blocking on all known confounders
- random treatment assignment within blocks
- $\Delta_{S_X} = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $\Delta_{T_X} = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small n
- little or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$
- $E(\Delta_{T_X}) = 0$
- $E(\Delta_{T_U}) = 0$

Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small n
- Full blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$
- $\Delta_{T_X} = 0$
- $E(\Delta_{T_U}) = 0$

Social Science Field Experiment

- nonrandom selection
- large n
- limited or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$ or change PATE to SATE and $\Delta_{S_X} = 0$
- $\Delta_{S_U} \neq 0$ or change PATE to SATE and $\Delta_{S_U} = 0$
- $E(\Delta_{T_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_X}) = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

Survey Experiment

- random selection
- large n
- limited or no blocking
- random treatment assignment (only question wording treatments)
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $E(\Delta_{T_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_X}) = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

Observational Study, well-matched

- nonrandom selection
- large n
- no blocking
- nonrandom treatment assignment
- $\Delta_{S_X} \approx 0$ if representative, corrected by weighting, or for estimating SATE; or $\neq 0$ otherwise
- $\Delta_{S_U} \neq 0$
- $\Delta_{T_X} \approx 0$ (due to matching well)
- $\Delta_{T_U} \neq 0$ except by assumption


What is the Best Design?

- The ideal design is rarely feasible
- the Achilles heel of experimental studies: Δ_S and small n
- the Achilles heel of observational studies: Δ_U
- Each design accommodates best to the applications for which it was designed
- Effort in experimental studies: random assignment
- Effort in observational studies: measuring and adjusting for X (via matching or modeling)

Fallacies in Experimental Research

- Failure to block on all known covariates
 - seen incorrectly as requiring fewer assumptions (about what to block on)
 - In fact, blocking helps compared to not blocking (except in strange situations)
 - Blocking on relevant covariates is better, so choose carefully.
 - “Block what you can and randomize what you cannot” (Box et al.)
- Conducting t-tests to check for balance after random treatment assignment
 - variables blocked on balance exactly after treatment assignment; so if you're checking, you missed an opportunity to increase efficiency
 - if variables became available after treatment assignment, a t-test only checks for whether randomization was done appropriately
 - randomization balances on average; any one random assignment is not balanced exactly (which is why its better to block)

The Balance Test Fallacy in Matching Research



Randomly dropping observations “reduces” imbalance???

The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
 - Suppose $E(Y | T, X) = \theta + T\beta + X\gamma$
 - Bias in coefficient on T from regressing Y on T (without X):
 $E(\hat{\beta} - \beta | T, X) = G\gamma$ (where G are coefficients from a regression X on a constant and T)
 - Imbalance is due to G only.
 - If $G = 0$, bias=0
 - If $G \neq 0$, bias can be any size due to γ
 - If you cannot control G (and we don't usually try to minimize γ so we don't cook the books), then G must be reduced without limit.
- No threshold level is safe.