

Section 1: Probability Review

Molly Roberts

January 30, 2013

Outline

- 1 Basic Probability
- 2 Random Variables & Probability Distributions
- 3 Simulation
 - Example 1: Finding a Mean
 - Example 2: Probability
 - Example 3: Integrating the Normal Density

Three Axioms of Probability

Let S be the sample space and A be an event in S .

- 1 For any event A , $P(A) \geq 0$.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

- 1 For any event A , $P(A) \geq 0$.
e.g. The probability of getting a heads cannot be -1.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

- ① For any event A , $P(A) \geq 0$.
e.g. The probability of getting a heads cannot be -1.
- ② $P(S) = 1$.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

- ① For any event A , $P(A) \geq 0$.
e.g. The probability of getting a heads cannot be -1.
- ② $P(S) = 1$.
e.g. $P(\text{heads}) + P(\text{tails}) = 1$

Three Axioms of Probability

Let S be the sample space and A be an event in S .

- ① For any event A , $P(A) \geq 0$.
e.g. The probability of getting a heads cannot be -1.
- ② $P(S) = 1$.
e.g. $P(\text{heads}) + P(\text{tails}) = 1$
- ③ If $A_1, A_2, \dots, A_n$ are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

Three Axioms of Probability

Let S be the sample space and A be an event in S .

- ① For any event A , $P(A) \geq 0$.
e.g. The probability of getting a heads cannot be -1.
- ② $P(S) = 1$.
e.g. $P(\text{heads}) + P(\text{tails}) = 1$
- ③ If $A_1, A_2, \dots, A_n$ are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

e.g. $P(\text{rolling a 3 or rolling a 2}) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$

Conditional Probability


Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$




Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$




Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$



Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$



Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(A|B)P(B)$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(A|B)P(B)$$

$$P(B \cap A) = P(B|A)P(A)$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(A|B)P(B)$$

$$P(B \cap A) = P(B|A)P(A)$$

$$P(B \cap A) = P(A \cap B)$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(A|B)P(B)$$

$$P(B \cap A) = P(B|A)P(A)$$

$$P(B \cap A) = P(A \cap B)$$

$$P(A|B)P(B) = P(B|A)P(A)$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(A|B)P(B)$$

$$P(B \cap A) = P(B|A)P(A)$$

$$P(B \cap A) = P(A \cap B)$$

$$P(A|B)P(B) = P(B|A)P(A)$$

Bayes' Rule:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Side Note: Bayes Rule

Bayes rule will be extremely important going forward.

Why?

- Often we want to know $P(\theta|\text{data})$.
- But what we *do* know is $P(\text{data}|\theta)$.
- We'll be able to *infer* θ by using a variant of Bayes rule. Stay tuned.

Bayes' Rule:

$$P(\theta|y) = \frac{P(y|\theta)P(\theta)}{P(y)}$$

Outline

- 1 Basic Probability
- 2 Random Variables & Probability Distributions**
- 3 Simulation
 - Example 1: Finding a Mean
 - Example 2: Probability
 - Example 3: Integrating the Normal Density

Random Variables

Random Variables

A **random variable** is a function from S , the sample space, to $\mathbb{R}$ the real line,

Random Variables

A **random variable** is a function from S , the sample space, to $\mathbb{R}$ the real line, in other words a numerical value calculated from the outcome of a random experiment.

Random Variables

A **random variable** is a function from S , the sample space, to $\mathbb{R}$ the real line, in other words a numerical value calculated from the outcome of a random experiment.

e.g. When rolling a two dice, we may be interested in whether or not the sum of the two dice is 7.

Random Variables

A **random variable** is a function from S , the sample space, to $\mathbb{R}$ the real line, in other words a numerical value calculated from the outcome of a random experiment.

e.g. When rolling a two dice, we may be interested in whether or not the sum of the two dice is 7. Or we might be interested in the actual sum of the two dice.



Ex. Waiting for the Redline – How long will it take for the next T to get here?



Ex. Waiting for the Redline – How long will it take for the next T to get here?

$$X = \begin{cases} 1 & \text{if the redline arrives within 1 minute} \\ 2 & \text{if 1 – 2 minutes} \\ 3 & \text{if 2 – 3 minutes} \\ 4 & \text{if 3 – 4 minutes} \\ \vdots & \vdots \end{cases}$$

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

$$\Pr(X = 1) = \pi = .2.$$

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

$$\Pr(X = 1) = \pi = .2.$$

What's $\Pr(X=2)$?

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

$$\Pr(X = 1) = \pi = .2.$$

What's $\Pr(X=2)$?

$$\Pr(X = 2) = (1 - \pi)\pi = .8 \cdot .2 = .16.$$

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

$$\Pr(X = 1) = \pi = .2.$$

What's $\Pr(X=2)$?

$$\Pr(X = 2) = (1 - \pi)\pi = .8 \cdot .2 = .16.$$

What's $\Pr(X=3)$?

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

$$\Pr(X = 1) = \pi = .2.$$

What's $\Pr(X=2)$?

$$\Pr(X = 2) = (1 - \pi)\pi = .8 \cdot .2 = .16.$$

What's $\Pr(X=3)$?

$$\Pr(X = 3) = (1 - \pi)^2\pi = .8^2 \cdot .2 = .128$$

Ex. Waiting for the Redline Cont

Now, suppose the probability that the T comes in any given minute is a constant $\pi = .2$, and whether the T comes is independent of what has happened in previous periods.

What's $\Pr(X=1)$?

$$\Pr(X = 1) = \pi = .2.$$

What's $\Pr(X=2)$?

$$\Pr(X = 2) = (1 - \pi)\pi = .8 \cdot .2 = .16.$$


What's $\Pr(X=3)$?

$$\Pr(X = 3) = (1 - \pi)^2\pi = .8^2 \cdot .2 = .128$$

And generally...

$$\Pr(X = x) = (1 - \pi)^{x-1}\pi = .8^{x-1} \cdot .2$$

A Pictorial Representation of the PMF




X is a Geometric random variable with parameter $\pi = .2$.

Probability Density Function

Ex. Waiting for the Redline: an alternative model where Y is exact time the T arrives.


$$f(y) = \lambda e^{-\lambda y} = .25e^{-.25y}$$



Probability Density Function

Ex. Waiting for the Redline: an alternative model where Y is exact time the T arrives.

$$f(y) = \lambda e^{-\lambda y} = .25e^{-.25y}$$



Y is an Exponential random variable with parameter $\lambda = .25$.

Ex. Calculating probabilities for continuous RVs

Recall that to identify probabilities for continuous random variables we have to use:


$$P(Y \in A) = \int_A f(y)dy.$$

$$Pr(2 \leq y \leq 10) =$$

Probability Density Function

Ex. Waiting for the Redline.: an alternative model where Y is exact time the T arrives.


$$f(y) = \lambda e^{-\lambda y} = .25e^{-.25y}$$



Probability Density Function

Ex. Waiting for the Redline.: an alternative model where Y is exact time the T arrives.

$$f(y) = \lambda e^{-\lambda y} = .25e^{-.25y}$$



Ex. Calculating probabilities for continuous RVs

Recall that to identify probabilities we must use¹:

$$P(Y \in A) = \int_A f(y) dy.$$

$$Pr(2 \leq y \leq 10) = \int_2^{10} .25e^{-.25y} dy$$

¹Here we also use the fact that the antiderivative of e^{bx} is $\frac{e^{bx}}{b}$ 

Ex. Calculating probabilities for continuous RVs

Recall that to identify probabilities we must use¹:

$$P(Y \in A) = \int_A f(y) dy.$$

$$\begin{aligned} Pr(2 \leq y \leq 10) &= \int_2^{10} .25e^{-.25y} dy \\ &= -e^{-.25y} \Big|_2^{10} \\ &= -e^{-.25 \cdot 10} + e^{-.25 \cdot 2} \\ &\approx .525 \end{aligned}$$

¹Here we also use the fact that the antiderivative of e^{bx} is $\frac{e^{bx}}{b}$ 

Characteristics of all PDFs and PMFs:

Characteristics of all PDFs and PMFs:

1. The support is all y 's where $P(Y = y) > 0$.

Characteristics of all PDFs and PMFs:

1. The support is all y 's where $P(Y = y) > 0$.
2. Probability mass/density must integrate to 1

Characteristics of all PDFs and PMFs:

1. The support is all y 's where $P(Y = y) > 0$.
2. Probability mass/density must integrate to 1
3. $P(Y = y) \geq 0$ for all Y

Characteristics of all PDFs and PMFs:

1. The support is all y 's where $P(Y = y) > 0$.
2. Probability mass/density must integrate to 1
3. $P(Y = y) \geq 0$ for all Y

Characteristics of all PDFs and PMFs:

1. The support is all y 's where $P(Y = y) > 0$.
2. Probability mass/density must integrate to 1
3. $P(Y = y) \geq 0$ for all Y

Ex.:

1. $\int_0^{\infty} .25e^{-.25y} dy = -e^{-.25y}|_0^{\infty} = 0 + 1 = 1$
2. $.25e^{-.25y} \geq 0$ for all $y \in (0, \infty)$

Expectation

Expectation

Discrete Case:

Expectation

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where $P(X = x)$ is the probability mass function (PMF).

Expectation

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where $P(X = x)$ is the probability mass function (PMF).

Continuous Case:

Expectation

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where $P(X = x)$ is the probability mass function (PMF).

Continuous Case:

$$E(Y) = \int_{-\infty}^{\infty} yf(y)dy$$

where $f(y)$ is the probability density function (PDF).

Our running examples...

Discrete Time:

Our running examples...

Discrete Time:

$$\begin{aligned}
 E(X) &= \sum_i x_i P(X = x_i) \\
 &= \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2 \\
 &= 5
 \end{aligned}$$

Our running examples...

Discrete Time:

$$\begin{aligned}
 E(X) &= \sum_i x_i P(X = x_i) \\
 &= \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2 \\
 &= 5
 \end{aligned}$$

Continuous Time:

Our running examples...

Discrete Time:

$$\begin{aligned}
 E(X) &= \sum_i x_i P(X = x_i) \\
 &= \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2 \\
 &= 5
 \end{aligned}$$

Continuous Time:

$$\begin{aligned}
 E(Y) &= \int_{-\infty}^{\infty} y f(y) dy \\
 &= \int_0^{\infty} y \cdot .25 e^{-.25y} dy \\
 &= 4
 \end{aligned}$$

Expectation of a Function of a Random Variable

Now let's complicate things: suppose we want to find $E[g(X)]$, where $g(X)$ is any function of X .

Expectation of a Function of a Random Variable

Now let's complicate things: suppose we want to find $E[g(X)]$, where $g(X)$ is any function of X .

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables

Expectation of a Function of a Random Variable

Now let's complicate things: suppose we want to find $E[g(X)]$, where $g(X)$ is any function of X .

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables and

$$E[g(y)] = \int_{-\infty}^{\infty} g(y)f(y)dy$$

for continuous random variables.

Return of our running examples

Suppose we want to find $E[g(X)]$, where $g(X) = \sqrt{1+x}$.

Return of our running examples

Suppose we want to find $E[g(X)]$, where $g(X) = \sqrt{1+x}$.

$$\begin{aligned} E[g(X)] &= \sum_i g(x_i)P(X = x_i) \\ &= \sum_{x=1}^{\infty} \sqrt{1+x}(1-.2)^{x-1} \cdot .2 \end{aligned}$$

Return of our running examples

Suppose we want to find $E[g(X)]$, where $g(X) = \sqrt{1+x}$.

$$\begin{aligned} E[g(X)] &= \sum_i g(x_i)P(X = x_i) \\ &= \sum_{x=1}^{\infty} \sqrt{1+x}(1-.2)^{x-1} \cdot .2 \end{aligned}$$

$$\begin{aligned} E[g(Y)] &= \int_{-\infty}^{\infty} g(y)f(y)dy \\ &= \int_0^{\infty} \sqrt{1+y}.25e^{-.25y} dy \end{aligned}$$

Outline

- 1 Basic Probability
- 2 Random Variables & Probability Distributions
- 3 Simulation**
 - Example 1: Finding a Mean
 - Example 2: Probability
 - Example 3: Integrating the Normal Density

The Monte Carlo Method

Basic idea: rather than calculate quantities analytically using deterministic formulae, approximate quantities using random sampling.

The Monte Carlo Method

Basic idea: rather than calculate quantities analytically using deterministic formulae, approximate quantities using random sampling.



Simulating an expectation

$$E(X) = \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2$$

Simulating an expectation

$$E(X) = \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2$$

```
> draws <- rgeom(n = 100000, prob = .2)
```

Simulating an expectation

$$E(X) = \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2$$

```
> draws <- rgeom(n = 100000, prob = .2)
> mean(draws)
[1] 4.99796
```

Simulating an expectation

$$E(X) = \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2$$

```
> draws <- rgeom(n = 100000, prob = .2)
> mean(draws)
[1] 4.99796
```

$$E(Y) = \int_0^{\infty} y \cdot .25e^{-.25y} dy$$

Simulating an expectation

$$E(X) = \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2$$

```
> draws <- rgeom(n = 100000, prob = .2)
> mean(draws)
[1] 4.99796
```

$$E(Y) = \int_0^{\infty} y \cdot .25e^{-.25y} dy$$

```
> draws <- rexp(n = 100000, rate = .25)
```

Simulating an expectation

$$E(X) = \sum_{x_i=1}^{\infty} x_i (1 - .2)^{x_i-1} \cdot .2$$

```
> draws <- rgeom(n = 100000, prob = .2)
> mean(draws)
[1] 4.99796
```

$$E(Y) = \int_0^{\infty} y \cdot .25e^{-.25y} dy$$

```
> draws <- rexp(n = 100000, rate = .25)
> mean(draws)
[1] 4.008509
```

Neither of these are perfectly accurate but they become arbitrarily

Monte Carlo Integration

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)f(x)dx$$

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)f(x)dx$$

Monte Carlo Integration allows us to approximate it by simulating M values from $f(x)$ and calculating:

$$\hat{I}_M = \frac{1}{M} \sum_{i=1}^M g(x^{(i)})$$

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)f(x)dx$$

Monte Carlo Integration allows us to approximate it by simulating M values from $f(x)$ and calculating:

$$\hat{I}_M = \frac{1}{M} \sum_{i=1}^M g(x^{(i)})$$

By the Strong Law of Large Numbers, our estimate $\hat{I}_M$ is a simulation consistent estimator of I as $M \rightarrow \infty$ (our estimate gets better as we increase the number of simulations).

Back to our examples

$$E[g(X)] = \sum_{x=1}^{\infty} \sqrt{1+x}(1-.2)^{x-1}.2$$

Back to our examples

$$E[g(X)] = \sum_{x=1}^{\infty} \sqrt{1+x}(1-.2)^{x-1}.2$$

Approach:

1. Draw $M = 100000$ samples from the geometric distribution.

Back to our examples

$$E[g(X)] = \sum_{x=1}^{\infty} \sqrt{1+x} (1-.2)^{x-1} .2$$

Approach:

1. Draw $M = 100000$ samples from the geometric distribution.
2. Calculate $g(x^{(i)})$ for each.

Back to our examples

$$E[g(X)] = \sum_{x=1}^{\infty} \sqrt{1+x}(1-.2)^{x-1}.2$$

Approach:

1. Draw $M = 100000$ samples from the geometric distribution.
2. Calculate $g(x^{(i)})$ for each.
3. Find the mean of these.

```
> draws <- rgeom(n = 100000, prob = .2)
```

Back to our examples

$$E[g(X)] = \sum_{x=1}^{\infty} \sqrt{1+x}(1-.2)^{x-1}.2$$

Approach:

1. Draw $M = 100000$ samples from the geometric distribution.
2. Calculate $g(x^{(i)})$ for each.
3. Find the mean of these.

```
> draws <- rgeom(n = 100000, prob = .2)
```

```
> g.draws <- sqrt(1 + draws)
```

```
> mean(g.draws)
```

```
[1] 2.312
```

Back to our examples


$$E[g(X)] = \int_0^{\infty} \sqrt{1+y} \cdot 0.25e^{-.25y} dy$$

Approach:


1. Draw $M = 100000$ samples from the exponential distribution.
2. Calculate $g(x^{(i)})$ for each.
3. Find the mean of these.

```
> draws <- rexp(n = 100000, rate = .25)
> g.draws <- sqrt(1 + draws)
> mean(g.draws)
[1] 2.096
```

Simulation for Probability Problems




I have an urn composed of 5 red balls and 5 green balls. If I sample 4 balls without replacement from the urn, what is the probability of drawing 4 balls all of the same color?




Here are some steps:

1. Construct our population aka our urn.




Here are some steps:

1. Construct our population aka our urn.
2. Figure out how to take one sample from it.




Here are some steps:

1. Construct our population aka our urn.
2. Figure out how to take one sample from it.
3. Figure out a programming rule for determining whether our condition was met i.e. we drew "4 red balls or 4 green balls".



Here are some steps:

1. Construct our population aka our urn.
2. Figure out how to take one sample from it.
3. Figure out a programming rule for determining whether our condition was met i.e. we drew "4 red balls or 4 green balls".
4. Throw a for loop around it and sample repeatedly.



Here are some steps:

1. Construct our population aka our urn.
2. Figure out how to take one sample from it.
3. Figure out a programming rule for determining whether our condition was met i.e. we drew "4 red balls or 4 green balls".
4. Throw a for loop around it and sample repeatedly.
5. Determine what proportion of times our condition was successful.

1. Construct our population

Here are some possibilities:

```
> urn <- c("G","G","G","G","G","R","R","R","R","R")  
> urn  
[1] "G" "G" "G" "G" "G" "R" "R" "R" "R" "R"
```

1. Construct our population

Here are some possibilities:

```
> urn <- c("G","G","G","G","G","R","R","R","R","R")
> urn
[1] "G" "G" "G" "G" "G" "R" "R" "R" "R" "R"
```

```
> urn <- c(rep("red",5),rep("green",5))
> urn
[1] "red"    "red"    "red"    "red"    "red"
     "green" "green"  "green"  "green"  "green"
```

1. Construct our population

Here are some possibilities:

```
> urn <- c("G","G","G","G","G","R","R","R","R","R")
```

```
> urn
```

```
[1] "G" "G" "G" "G" "G" "R" "R" "R" "R" "R"
```

```
> urn <- c(rep("red",5),rep("green",5))
```

```
> urn
```

```
[1] "red"    "red"    "red"    "red"    "red"
     "green" "green"  "green"  "green"  "green"
```

```
> urn <- c(rep(1,5),rep(0,5))
```

```
> urn
```

```
[1] 1 1 1 1 1 0 0 0 0 0
```

1. Construct our population

Here are some possibilities:

```
> urn <- c("G","G","G","G","G","R","R","R","R","R")
> urn
[1] "G" "G" "G" "G" "G" "R" "R" "R" "R" "R"
```

```
> urn <- c(rep("red",5),rep("green",5))
> urn
[1] "red"    "red"    "red"    "red"    "red"
     "green" "green"  "green"  "green"  "green"
```

```
> urn <- c(rep(1,5),rep(0,5))
> urn
[1] 1 1 1 1 1 0 0 0 0 0
```

I'll use this last one because numbers will be easier to use later on.

2. Figure out how to take one sample

We need to use the `sample()` function with the `replace = FALSE` argument:

```
> sample(x = urn, size = 4, replace = FALSE)
[1] 1 1 1 0
```

3. Determine if a success or failure

Because we used numeric signifiers for red and green, there is an easy test for whether or not we have drawn balls of all one color. If the numbers sum up to either 0 or 4, then we have a 'success'.

3. Determine if a success or failure

Because we used numeric signifiers for red and green, there is an easy test for whether or not we have drawn balls of all one color. If the numbers sum up to either 0 or 4, then we have a 'success'.

```
> draw <- sample(x = urn, size = 4, replace = FALSE)
> draw
[1] 0 0 0 0
```

3. Determine if a success or failure

Because we used numeric signifiers for red and green, there is an easy test for whether or not we have drawn balls of all one color. If the numbers sum up to either 0 or 4, then we have a 'success'.

```
> draw <- sample(x = urn, size = 4, replace = FALSE)
> draw
[1] 0 0 0 0

> sum(draw) == 4
[1] FALSE

> sum(draw) == 0
[1] TRUE
```

3. Determine if a success or failure

Because we used numeric signifiers for red and green, there is an easy test for whether or not we have drawn balls of all one color. If the numbers sum up to either 0 or 4, then we have a 'success'.

```
> draw <- sample(x = urn, size = 4, replace = FALSE)
> draw
[1] 0 0 0 0

> sum(draw) == 4
[1] FALSE

> sum(draw) == 0
[1] TRUE
```

But we can combined these test using '|' which means 'or'

```
> (sum(draw) == 4) | (sum(draw) == 0)
[1] TRUE
```

4. Throw a for-loop around it

Here's our guts so far:

```
draw <- sample(x = urn, size = 4, replace = FALSE)
success <- sum(draw) == 4 | sum(draw) == 0
```

4. Throw a for-loop around it

Here's our guts so far:

```
draw <- sample(x = urn, size = 4, replace = FALSE)
success <- sum(draw) == 4 | sum(draw) == 0
```

We can repeat this over and over:

```
sims <- 100000
success <- NULL
for(i in 1:sims){
  draw <- sample(x = urn, size = 4, replace = FALSE)
  success[i] <- sum(draw) == 4 | sum(draw) == 0
}
```

(4. Throw replicate around it)

for-loops in R are slow. To get fancy:

```
eval <- function(draw){  
  draw <- sample(x = urn, size = 4, replace = FALSE)  
  success <- sum(draw) == 4 | sum(draw) == 0  
  return(success)  
}
```

(4. Throw replicate around it)

for-loops in R are slow. To get fancy:

```
eval <- function(draw){  
  draw <- sample(x = urn, size = 4, replace = FALSE)  
  success <- sum(draw) == 4 | sum(draw) == 0  
  return(success)  
}
```

We can use replicate to make the loop faster:

```
sims <- 100000  
success <- replicate(sims, eval(sample(urn, 4, replace=F)))
```

5. Determine proportion of success

Here are two equivalent approaches.

```
> sum(success)/sims
```

```
[1] 0.047
```

```
> mean(success)
```

```
[1] 0.047
```

Why is this useful?

- Math is hard or impossible and takes too long.

Why is this useful?

- Math is hard or impossible and takes too long.
- Consider trying to integrate

$$I(f) = \frac{1}{\sqrt{2\pi}} \int_0^1 e^{-\frac{x^2}{2}} dx$$

Why is this useful?

- Math is hard or impossible and takes too long.
- Consider trying to integrate

$$I(f) = \frac{1}{\sqrt{2\pi}} \int_0^1 e^{-\frac{x^2}{2}} dx$$

Which is the standard normal density and cannot be evaluated in closed form.

Why is this useful?

- Math is hard or impossible and takes too long.
- Consider trying to integrate

$$I(f) = \frac{1}{\sqrt{2\pi}} \int_0^1 e^{-\frac{x^2}{2}} dx$$

Which is the standard normal density and cannot be evaluated in closed form.

- How could we solve this?

Why is this useful?

- Math is hard or impossible and takes too long.
- Consider trying to integrate

$$I(f) = \frac{1}{\sqrt{2\pi}} \int_0^1 e^{-\frac{x^2}{2}} dx$$

Which is the standard normal density and cannot be evaluated in closed form.

- How could we solve this?
 - 1 Sample 1000 points, $X_1, \dots, X_{1000}$, uniformly distributed over the interval $(0, 1)$.

Why is this useful?

- Math is hard or impossible and takes too long.
- Consider trying to integrate

$$I(f) = \frac{1}{\sqrt{2\pi}} \int_0^1 e^{-\frac{x^2}{2}} dx$$

Which is the standard normal density and cannot be evaluated in closed form.

- How could we solve this?
 - ① Sample 1000 points, $X_1, \dots, X_{1000}$, uniformly distributed over the interval $(0, 1)$.
 - ② Evaluate the function at each of these points and take the mean.

$$\frac{1}{1000} \left(\frac{1}{\sqrt{2\pi}} \right) \sum_{i=1}^{1000} e^{-\frac{x_i^2}{2}}$$